


<http://mc.manuscriptcentral.com/fems>

**Environmental dimensions of antibiotic resistance:  
Assessment of basic science gaps**

Journal:	<i>FEMS Microbiology Ecology</i>
Manuscript ID	FEMSEC-18-09-0521
Manuscript Type:	Perspective
Date Submitted by the Author:	21-Sep-2018
Complete List of Authors:	Smalla, Kornelia; Julius Kühn-Institute, Federal Research Centre for Cultivated Plants, Institute for Epidemiology and Pathogen Diagnostics Cook, Kimberly; USDA_ARC, Micro Djordjevic, Steven; University of Technology, Sydney, Professor, The itthree Institute Klümper, Uli; University of Exeter, Environment and Sustainability Institute Gillings, Michael; Macquarie University, Biological Sciences;
Keywords:	plasmid, insertion element, microbiome, One health, evolution, pathogen

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



190x254mm (96 x 96 DPI)

1 **Perspective – FEMS Microbial Ecology**

2 **EDAR 4 Roundtable 1**

3 **Environmental dimensions of antibiotic resistance: Assessment of basic science gaps**

4  
5 Kornelia Smalla<sup>1</sup>, Kimberley Cook<sup>2</sup>, Steven P. Djordjevic<sup>3</sup>, Uli Klümper<sup>4</sup>, and Michael  
6 Gillings<sup>5</sup>

7  
8 <sup>1</sup>Julius Kühn-Institut Federal Research Centre for Cultivated Plants, Institute for  
9 Epidemiology and Pathogen Diagnostics, Braunschweig, Germany, <sup>2</sup>Bacterial Epidemiology  
10 and Antimicrobial Resistance Research Unit, U.S. National Poultry Research Center, USDA  
11 Agricultural Research Service, Athens, Georgia, USA, <sup>3</sup>ithree institute, University of  
12 Technology Sydney, Broadway, Sydney, NSW, Australia, <sup>4</sup>ESI & CEC, Biosciences,  
13 University of Exeter, Penryn Campus, Cornwall, UK, <sup>5</sup>Department of Biological Sciences,  
14 Macquarie University, Sydney, NSW, Australia.

15  
16 Correspondence:

17 Prof Michael Gillings

18 Biological Sciences

19 Macquarie University

20 NSW 2019, Australia

21 [michael.gillings@mq.edu.au](mailto:michael.gillings@mq.edu.au)

22 61 2 9850 8199

23  
24 Keywords: plasmid; insertion element; microbiome; One Health; evolution; pathogen

**Abstract**

Antibiotic resistance is one of the major problems facing medical practice in the 21<sup>st</sup> century. Historical approaches to managing antibiotic resistance have often focused on individual patients, specific pathogens, and particular resistance phenotypes. However, it is increasingly recognized that antibiotic resistance is a complex ecological and evolutionary problem. As such, understanding the dynamics of antibiotic resistance requires integration of data on the diverse mobile genetic elements often associated to antibiotic resistance genes, and their dissemination through various mechanisms of horizontal gene transfer between bacterial cells and environments. Most important is understanding of the fate and effects of antibiotics at sub-inhibitory concentration and co-selection. This opinion paper identifies key knowledge gaps in our understanding of resistance phenomena, and outlines research needs that should be addressed to help us manage resistance into the future.

## 37 Introduction

38 Over the past 80 years, antibiotics have revolutionised modern medicine because of their  
39 ability to control bacterial infections. However, the use of antibiotics in medicine, and during  
40 animal and plant production, imposes strong selection pressures on microbial communities  
41 exposed to sub-inhibitory concentration of antibiotics. This has driven the rise of resistance,  
42 via mutation and horizontal gene transfer of antibiotic resistance genes (ARGs) in diverse  
43 habitats and most prominently in so-called hot spots of horizontal gene transfer (Davies and  
44 Davies, 2010; Heuer and Smalla, 2012; Gillings, 2017).

45 The rise of resistance to all classes of antibiotics poses a major threat to modern health  
46 practice. Understanding and controlling the ongoing dissemination of antibiotic resistant  
47 bacteria is arguably one of the most pressing tasks for managing human health in the 21<sup>st</sup>  
48 century (Laxminarayan *et al.*, 2013). Like many other pressing human problems,  
49 investigating antibiotic resistance involves examining complex, dynamic processes that  
50 reflect the global changes in microbial communities induced by humanity (Zhu *et al.*, 2017a).

51 Mobile genetic elements (MGEs) are an important component of the flexible gene pool of  
52 bacterial cells allowing bacterial communities to rapidly adapt to changing environmental  
53 conditions e.g. through the presence of antibiotics (Heuer and Smalla, 2012). There is a  
54 pressing need to gather comprehensive knowledge about the diversity and dissemination of  
55 MGEs that often carry various resistance genes and thus foster their environmental spread.  
56 Characterizing complete plasmid sequences and understanding the mechanisms that generate  
57 mobile DNAs should be a priority (Chowdhury *et al.*, 2015; San Millan *et al.*, 2016;  
58 McKinnon *et al.*, 2018). Understanding the transfer of DNA elements from the environmental  
59 resistome into pathogens and commensals is also critical (D'Costa *et al.*, 2006; Gaze *et al.*,  
60 2013; Wellington *et al.*, 2013). By identifying the physical locations and genetic systems  
61 where the most rapid change is occurring, we might be able to intervene, and identify  
62 potential control points where mitigation strategies might be most effective. However, in  
63 many areas, we still lack fundamental information about the origins, dissemination and  
64 maintenance of ARGs. To understand these dynamics, we will need to integrate complex data  
65 sets that span different biological, temporal and spatial scales and link these with abiotic and  
66 biotic environmental factors.

67 Most importantly, understanding and mitigating the dissemination of antibiotic resistance  
68 requires a better understanding of the ecology of MGEs, ARGs and their hosts. Each habitat

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

69 is populated by complex and dynamic bacterial communities that differ in their taxonomic  
70 composition. These differences in taxonomic composition influence the type and prevalence  
71 of MGEs such as plasmids, integrated conjugative elements, phages, transposons and  
72 integrons. Land use practices and anthropogenic pollutants in waste streams influence the  
73 composition of bacterial communities, potentially selecting lineages that carry adaptive traits  
74 and affecting microbial diversity.

75 In the roundtable discussion at EDAR-4, we identified some key knowledge gaps (Figure 1),  
76 and suggested research needs and paths forward to resolving these difficulties.

77  
78 **Gaps in our understanding of baselines and microbial ecology**

79 The modern world is a consequence of historical contingencies (Lowenthal, 2015). We live in  
80 an environment where the abundance, distribution and activities of microbial communities,  
81 their genes, and the MGEs they carry have been profoundly altered through anthropogenic  
82 activity over the last 50 years (Gillings and Paulsen, 2014; Zhu *et al.*, 2017b). This means  
83 that we potentially have no baseline of unaffected or pristine samples from some  
84 environments with which we can make comparisons.

85 The dissemination routes of bacteria that carry MGEs and the persistence and mobilization of  
86 MGE are not well understood. The relative contributions of transport mechanisms such as of  
87 water, wind, dust, migrating wildlife or human activities (tourism, waste management,  
88 agriculture) are unclear. In turn, this makes it difficult to estimate rates of increase in  
89 abundance and prevalence, especially when some environmental compartments have  
90 potentially reached saturated concentrations of antibiotics, metal compounds and ARGs. It  
91 has been assumed that the relative abundance of bacteria carrying transferable ARGs is  
92 higher in the presence of pollutants (Jechalke *et al.*, 2014), since under these circumstances,  
93 plasmid carriage presents a fitness advantage. Certainly, in the absence of such selective  
94 pressures, the proportion of the population that carries antibiotic resistance plasmids declines  
95 (Jechalke *et al.*, 2013).

96 In theory, because of the costs of replication and maintenance, plasmids should be lost from  
97 their hosts in the long-term due to genome streamlining, regardless of their short-term fitness  
98 benefits. Complete plasmid loss occurs when plasmid-free daughter cells outcompete cells  
99 that still carry the plasmid. This phenomenon can occur even when under antibiotic selection

100 pressure if the beneficial resistance gene becomes integrated into the chromosome  
101 (Bergstrom *et al.*, 2000; Harrison *et al.*, 2015). However, it was also proposed that plasmid  
102 carrying bacterial cells might present only a small proportion of a respective population  
103 (multi-cellular behaviour) and that the proportion of these plasmid carrying cells is increased  
104 under conditions where plasmid carriage represents a selective advantage (Heuer and Smalla,  
105 2012). Alternatively, compensatory mutations can occur after a plasmid is introduced into a  
106 new host, reducing the metabolic burden of plasmid maintenance (Harrison *et al.*, 2016), or  
107 turning the cost into a benefit (Loftie-Eaton *et al.*, 2017). These compensatory mutations can  
108 increase long-term retention of plasmids. Broad host range plasmids can then spread to  
109 diverse new hosts (Klümper *et al.*, 2014, 2017) in which similar co-evolution can occur.

110 Consequently, it is difficult to estimate the half-lives of resistant bacteria and their cargoes of  
111 mobile ARGs because it depends on the particular bacterial species, plasmids, phenotypic  
112 traits and environments involved. Knowing the rates of gain and loss in different  
113 environmental compartments is essential to assess the effectiveness of mitigation strategies  
114 (Figure 1).

115 There are, however, a few studies that have given us an idea of baselines in the pre-antibiotic  
116 era. Examination of resistance genes in archived soil samples (Knapp *et al.*, 2010), samples  
117 from permafrost sediments (D'Costa *et al.*, 2011), or in historical plasmid collections  
118 (Hughes and Datta, 1983), has given us a glimpse of this pre-antibiotic world. These studies  
119 suggest that clinically relevant resistance genes were rare in soils, and were not found on  
120 plasmids, prior to the 1940s. We suggest that there should be a concerted effort to examine  
121 archived samples, such as soils, seeds, pathology specimens, herbaria and culture collections  
122 from before the antibiotic era. This would help us distinguish the naturally occurring  
123 resistome (Allen *et al.*, 2010) from those resistome elements whose abundance has markedly  
124 increased under human influence (Zhu *et al.*, 2017b, 2017a).

125 We do know that agricultural practices alter the soil microbiome (Degrunen *et al.*, 2017;  
126 Hartmann *et al.*, 2017) and thus most likely also the relative abundance of resistance genes  
127 and mobile genetic elements in soil. However, the effect of agricultural management on the  
128 abundance and diversity of ARGs and MGEs is a clear knowledge gap. In general, those soils  
129 that have been exposed to organic fertilizers such as manure, digestates or sewage sludge  
130 have readily detectable resistance genes and mobile genetic elements, while these DNAs are  
131 below the detection limit in soils without previous history of organic fertilizer application

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

(Blau *et al.*, 2018b; Wolters *et al.*, 2018). Organic fertilizers are not only a valuable source of nutrients for plant growth but they contain also antibiotics and are a reservoir of ARGs and MGEs (Zhu *et al.*, 2013; Jechalke *et al.*, 2014; Wolters *et al.*, 2016). Furthermore, soil communities that have been subject to manure application display a significantly increased permissiveness towards mobile genetic elements, such as plasmids encoding antimicrobial resistance (Musovic *et al.*, 2014).

Antibiotic residues in manure applied to soil can influence selection and maintenance of resistance genes. This effect will depend strongly on the half-life of the antibiotic, and its bioavailability, which in turn will be dependent on soil type. However, even sub-lethal concentrations of antibiotics are known to select for cells carrying resistance plasmids encoding selective traits (Gullberg *et al.*, 2011, 2014) but might also select for mutations. Multiple mutations in genes that are not associated with resistance can result in highly resistant phenotypes (Wistrand-Yuen *et al.*, 2018). In general, the effects of antibiotics will be further compounded by potential co-selection by contaminating metal compounds, quaternary ammonium compounds and other biocides (Baker-Austin *et al.*, 2006; Jechalke *et al.*, 2014; Song *et al.*, 2017). In this regard, it is worth noting that diverse, multiple drug resistant commensal *E. coli* from pigs can also carry various heavy metal resistance genes (Reid *et al.*, 2017). Understanding the fate of antibiotics in different environmental settings, their interaction with other agents, and their role in selection within environmental compartments is a major gap in our understanding of resistance dynamics (Figure 1).

Bacteria that carry ARGs can occur on the surface of foods that are eaten raw, such as some vegetables. This represents a route of dissemination that deserves more attention, as it represents a direct link between the environment and humans. Transposons and integrons associated with resistance genes have been traced back to leafy vegetables, which have been suggested as the original source of these elements (Ghaly *et al.*, 2017). After incubation of lettuce or cilantro leaves in peptone broth, multiple resistance plasmids were detected in *E. coli* isolates, and in total community DNA obtained from the enrichment (Blau *et al.*, 2018a).

Ingesting resistant bacteria from the environment is directly correlated with a higher risk of gut colonization by these resistant phenotypes. Surfers, for example, are up to 4-fold more likely to be colonised by *E. coli* harbouring *bla*<sub>CTX-M</sub> genes compared to non-surfers, due to an increased ingestion of bathing waters contaminated with resistant bacteria (Leonard *et al.*, 2018). While these resistant bacteria might be lost after 2 to 6 months without exposure



(Kennedy and Collignon, 2010), regular intake of novel resistant bacteria via fresh produce could lead to a constant state of gut colonization. These bacteria can then be transported between continents as residents in the gut of tourists (Bengtsson-Palme *et al.*, 2015).

Based on these studies and others like them, it is clear that general conclusion can often not be drawn as multiple abiotic and biotic factors influence the findings and the ecologies of those systems need to be considered. Physical, chemical, spatial, temporal and biological complexities of natural systems and the ecologies of resistance in those systems preclude a “one size fits all” approach (Durso and Cook, 2014). Scales of environmental studies (i.e., laboratory, experimental/model, and spatial) must also be considered as part of hypothesis driven research. Identification of risk factors, bottlenecks and drivers of resistance may require different strategies for analysis and, ultimately, for remediation. Identifying and communicating applied solutions to critical communities in the field, including farmers, public health, regulatory groups and educators, represents a critical gap that limits implementation of solutions (Figure 1).

### **Gaps in understanding mechanisms of capture and dissemination**

Human attempts to control bacterial growth with antibiotics lead to selection of resistant bacterial lineages, and changes in bacterial community composition. However, a surprising amount of complexity and uncertainty lies behind this apparently obvious statement. What is the relative importance of mechanisms that influence resistance dynamics? Several mechanisms, such as conjugation, require proximity of donor and recipient cell. In other mechanisms, such as transformation, transduction, and DNA transfer by vesicles or gene transfer agents, donors and recipients can be separated temporally and spatially (Gillings, 2017). While conjugation is generally considered to be the most important mechanism, this may simply be because it is more tractable in laboratory and field experiments.

It is abundantly clear that the raw materials for selection of resistance are mutations to existing genes, or horizontal gene transfer events that can confer resistance phenotypes. However, what drives the initial horizontal gene transfer events, what mechanisms are important, where selection occurs, and the units of selection are less clear (Figure 1).

Most likely the importance of different mechanisms depends on the bacterial species, the genes, and the plasmids involved. However, there is also strong evidence that there are

1  
2  
3 195 environmental hot spots for resistance gene evolution and horizontal gene transfer (Rizzo *et*  
4 196 *al.*, 2013; Berendonk *et al.*, 2015). High cell densities, the availability of nutrients allowing  
5  
6 197 bacterial growth and the presence of pollutants are assumed to foster these processes. These  
7  
8 198 conditions can be found within the intestine of humans or animals treated with antibiotics, in  
9  
10 199 sewage treatment plants, animal manures, biofilters and in the rhizosphere. Biofilms are  
11  
12 200 typically also important sites that foster horizontal transfer of ARGs and MGEs (Flemming *et*  
13 201 *al.*, 2016; Gillings *et al.*, 2017).

14  
15 202 To fill the gaps in our limited understanding of the mechanisms of capture and dissemination  
16  
17 203 of ARGs the molecular characterization of MGEs and their ecology is of key importance.  
18  
19 204 Thus sequencing plasmids and complete bacterial genomes shows that plasmids from  
20  
21 205 unpolluted sites carry transposons, IS elements and integrons, but without ARGs. The same  
22  
23 206 plasmid types isolated from hospital environments, sewage sludge, manure or digestates do  
24  
25 207 carry ARGs. Consequently, plasmids appear to be an important means by which bacterial  
26  
27 208 hosts can dynamically and rapidly respond to changing environmental conditions by  
28  
29 209 acquiring advantageous phenotypes. Precisely where the acquisition of resistance genes takes  
30  
31 210 place, and what factors foster these processes remains unclear. While plasmid carriage under  
32  
33 211 conditions of rapid growth might be costly, this might not be the case under environmental  
34  
35 212 conditions. There is a gap in our understanding of horizontal transfer, metabolic costs of  
36  
37 213 plasmids and resistance gene capture under *in situ* conditions and thus research addressing  
38  
39 214 these topics needs to be done under more relevant conditions.

40  
41 215 Better understanding the role played by complex interactions between different mobile  
42  
43 216 elements is thus critical. This is perhaps best exemplified by the capture of chromosomally-  
44  
45 217 derived *bla*<sub>KLUA</sub> genes from *Kluyvera* species that inhabit the plant rhizosphere, onto  
46  
47 218 conjugative plasmids circulating widely in the Enterobacteriaceae. The *bla*<sub>KLUA</sub> genes have  
48  
49 219 close sequence identity to the *bla*<sub>CTX-M</sub> genes that are now globally disseminated in Gram-  
50  
51 220 negative bacterial populations from clinical environments. The insertion element *ISEcp1*  
52  
53 221 most likely played a key role in mobilising *bla*<sub>KLUV</sub> genes onto plasmids (Humeniuk *et al.*,  
54  
55 222 2002; Zhao and Hu, 2013).

56  
57 223 IS26 is another example for an insertion element that is shaping the accessory gene load of  
58  
59 224 drug resistance plasmids and the structure of class 1 integrons, promoting the formation of  
60  
225 clustered resistance gene regions (Cain *et al.*, 2010; Harmer and Hall, 2016; Reid *et al.*,  
226 2017). Copies of IS26 are known to flank diverse ARGs, creating independently mobile,

compound transposons. The association of IS26 with class 1 integrons shows how evolutionary events driven by constant selection pressure can assemble a powerful combination of disparate genetic elements with a formidable capacity to capture and express ARGs. Furthermore, IS26 can influence the formation and ongoing evolution of virulence plasmids via the creation of hybrid plasmids that merge virulence genes with ARGs (Venturini *et al.*, 2010; García *et al.*, 2016; Mangat *et al.*, 2017; Wong *et al.*, 2017). Perhaps of even greater concern is the role played by IS26-mediated deletion events in the promotion of plasmid fitness and host range. Newly-acquired plasmids may persist in a host background for long enough to enable IS-mediated modifications to partially alleviate the fitness imposed of carrying such plasmids (Porse *et al.*, 2016). IS26 is disseminated widely in commensal bacterial populations in the gastrointestinal tracts of food animals (Reid *et al.*, 2017), giving widespread opportunity for such events to occur. These studies are in their infancy and there is a pressing need to understand the role of commensal bacterial populations in the creation and spread of drug resistance elements. Experiments need to be designed to understand these events under *in situ* conditions.

There are still unaddressed issues of scale in the selection of resistant lineages. Selection effects can make themselves felt at scales of the cell membrane, within biofilms, soil particles, local populations, the vertebrate gut, or even at the landscape scale. Each scale has inherently different potentials for generating gradients of selective agents. The targets of selection are also nested at different scales, and include: the resistance gene; the mobile element it may reside upon; the genomic landscape around the resistance gene; the bacterial species involved; and the environmental compartments or the host that carries the bacterium (Figure 1). The intensity and direction of selection may differ across these scales, and it is a priority to tease out these complexities with empirical studies.

## Gaps and problems with information and data

It is fair to say that much of the energy devoted to investigating the environmental dimensions of antibiotic resistance has been in descriptive studies. The analysis of total community DNA directly extracted from environmental samples allowed comparative studies on the prevalence and relative abundance of ARGs and MGEs, and at the same time allowed characterization of microbial community composition. These studies have assembled large amounts of data about different sample types, different genes and different species.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Now it is time to pool these data, and this, in turn, will require standardization of nomenclature, metrics and annotation. However, it is also important to note that metagenomic studies, even with deep sequencing efforts, will likely only capture the resistome and mobilome of the most abundant bacteria. Furthermore, it is important to demonstrate functionality and transferability of the resistance genes detected. Therefore, multi-phasic approaches should be used that combine cultivation-dependent and independent methods.

While assembling and analysing these data, we must bear in mind that there is considerable errors and biases in those data that have already been collected. Some of this bias is a historical focus on clinically important species, and on those species amenable to genome sequencing. We need to understand the contribution of yet to be cultured organisms and of the rare microbiome, and this will require metagenomic analyses (Bengtsson-Palme *et al.*, 2017). However, depending on their genetic localization, resistance determinants might be shared by phylogenetically distant taxa. Any one gene can be found in multiple species, in diverse genetic contexts, and linked to transposons and IS elements that can occur in multiple copies in any one genome. This makes accurate assembly of metagenomic data challenging, especially for short read technologies.

**Education on microbial ecology, the natural role of antibiotics and how they affect the microbiota**

In the last decade numerous discoveries revealed the importance of a diverse microbiota for human well-being and for plant health. Antibiotic treatments disturb these communities and might affect their functionality. Thus prudent use of antibiotics should be an imperative. The rapid spread of ARGs demonstrates that bacteria dynamically respond and adapt to changing environments and stress. The most important message here is that the environmental dimension of antibiotic resistance is closely linked to the veterinary and human dimension of antibiotic resistance. Bacteria carrying ARGs and MGEs are exchanged through multiple paths between the environmental settings and thus the One Health perspective is a logical response to the resistance crisis (Robinson *et al.*, 2016; McEwen and Collignon, 2018).

It is important to note that the causal agents of nosocomial infections were often affiliated to genera such as *Acinetobacter*, *Burkholderia*, *Enterobacter*, *Klebsiella*, *Pseudomonas*, *Stenotrophomonas* or *Serratia* that are typically known as plant associated bacteria (Ryan *et al.*, 2009; García-León *et al.*, 2014). They have the ability to form biofilms and they display

multiple antibiotic resistances, and thus they have enormous selective advantages in the face of human attempts to control bacterial growth. These species potentially serve as important reservoirs of transferable ARGs. The rare plant microbiome also contains *E. coli* carrying transferable multiple resistance plasmids.

For too long, the problem of antibiotic resistance has been thought of as one of individual patients, infected with individual resistant organisms. It is now clear that the resistance crisis, is at its heart, a phenomenon of ecology and evolution. Educating patients, clinicians, farmers and regulators about the environmental dimensions of this problem is essential if we are to make progress. We need to acknowledge that while resistance determinants often originate from environmental compartments, these elements are now so common in human dominated ecosystems, that humans themselves have become a significant source of resistance pollution (Zhu *et al.*, 2017a; Gillings, 2018). Like many of the other environmental crises faced by humanity, antibiotic resistance is now a global problem that requires global solutions.

## References

- Allen HK, Donato J, Wang HH *et al.* Call of the wild: Antibiotic resistance genes in natural environments. *Nat Rev Microbiol* 2010;**8**:251–9.
- Baker-Austin C, Wright MS, Stepanauskas R *et al.* Co-selection of antibiotic and metal resistance. *Trends Microbiol* 2006;**14**:176–82.
- Bengtsson-Palme J, Angelin M, Huss M *et al.* The human gut microbiome as a transporter of antibiotic resistance genes between continents. *Antimicrob Agents Chemother* 2015;**59**:6551–60.
- Bengtsson-Palme J, Larsson DGJ, Kristiansson E. Using metagenomics to investigate human and environmental resistomes. *J Antimicrob Chemother* 2017;**72**:2690–703.
- Berendonk TU, Manaia CM, Merlin C *et al.* Tackling antibiotic resistance: the environmental framework. *Nat Rev Microbiol* 2015;**13**:310–7.
- Bergstrom CT, Lipsitch M, Levin BR. Natural selection, infectious transfer and the existence conditions for bacterial plasmids. *Genetics* 2000;**155**:1505–19.
- Blau K, Bettermann A, Jechalke S *et al.* The transferable resistome of produce. *MBio* 2018a. accepted:350629.
- Blau K, Casadevall L, Wolters B *et al.* Soil texture-depending effects of doxycycline and streptomycin applied with manure on the bacterial community composition and resistome. *FEMS Microbiol Ecol* 2018b;**94**:1–11.
- Cain AK, Liu X, Djordjevic SP *et al.* Transposons related to Tn1696 in IncHI2 plasmids in multiply antibiotic resistant *Salmonella enterica* serovar typhimurium from Australian animals. *Microb Drug Resist* 2010;**16**:197–202.

- 327 Chowdhury PR, Charles IG, Djordjevic SP. A role for Tn6029 in the evolution of the  
328 complex antibiotic resistance gene loci in genomic island 3 in enteroaggregative  
329 hemorrhagic escherichia coli O104:H4. *PLoS One* 2015;**10**:e0115781.
- 330 D'Costa VM, King CE, Kalan L *et al.* Antibiotic resistance is ancient. *Nature* 2011;**477**:457–  
331 61.
- 332 D'Costa VM, McGrann KM, Hughes DW *et al.* Sampling the antibiotic resistome. *Science*  
333 2006;**311**:374–7.
- 334 Davies J, Davies D. Origins and evolution of antibiotic resistance. *Microbiol Mol Biol Rev*  
335 2010;**74**:417–33.
- 336 Degruene F, Theodorakopoulos N, Colinet G *et al.* Temporal dynamics of soil microbial  
337 communities below the seedbed under two contrasting tillage regimes. *Front Microbiol*  
338 2017;**8**:1127.
- 339 Durso LM, Cook KL. Impacts of antibiotic use in agriculture: What are the benefits and  
340 risks? *Curr Opin Microbiol* 2014;**19**:37–44.
- 341 Flemming HC, Wingender J, Szewzyk U *et al.* Biofilms: An emergent form of bacterial life.  
342 *Nat Rev Microbiol* 2016;**14**:563–75.
- 343 García-León G, Hernández A, Hernando-Amado S *et al.* A function of SmeDEF, the major  
344 quinolone resistance determinant of *Stenotrophomonas maltophilia*, is the colonization  
345 of plant roots. *Appl Environ Microbiol* 2014;**80**:4559–65.
- 346 García V, García P, Rodríguez I *et al.* The role of IS26 in evolution of a derivative of the  
347 virulence plasmid of *Salmonella enterica* serovar Enteritidis which confers multiple  
348 drug resistance. *Infect Genet Evol* 2016;**45**:246–9.
- 349 Gaze WH, Krone SM, Joakim Larsson DG *et al.* Influence of humans on evolution and  
350 mobilization of environmental antibiotic resistome. *Emerg Infect Dis* 2013;**19**, DOI:  
351 10.3201/eid1907.120871.
- 352 Ghaly TM, Chow L, Asher AJ *et al.* Evolution of class 1 integrons: Mobilization and  
353 dispersal via food-borne bacteria. *PLoS One* 2017;**12**:e0179169.
- 354 Gillings MR. Lateral gene transfer, bacterial genome evolution, and the Anthropocene. *Ann N*  
355 *Y Acad Sci* 2017;**1389**:20–36.
- 356 Gillings MR. DNA as a pollutant: the clinical class 1 integron. *Curr Pollut Reports*  
357 2018;**4**:49–55.
- 358 Gillings MR, Paulsen IT. Microbiology of the anthropocene. *Anthropocene* 2014;**5**:1–8.
- 359 Gillings MR, Paulsen IT, Tetu SG. Genomics and the evolution of antibiotic resistance. *Ann*  
360 *N Y Acad Sci* 2017;**1388**:92–107.
- 361 Gullberg E, Albrecht LM, Karlsson C *et al.* Selection of a multidrug resistance plasmid by  
362 sublethal levels of antibiotics and heavy metals. *MBio* 2014;**5**:e01918-14-e01918-14.
- 363 Gullberg E, Cao S, Berg OG *et al.* Selection of resistant bacteria at very low antibiotic  
364 concentrations. *PLoS Pathog* 2011;**7**:e1002158.
- 365 Harmer CJ, Hall RM. IS 26 - mediated formation of transposons carrying antibiotic resistance  
366 genes. Blokesch M (ed.). *mSphere* 2016;**1**:e00038-16.



- 367 Harrison E, Dytham C, Hall JPJ *et al.* Rapid compensatory evolution promotes the survival  
368 of conjugative plasmids. *Mob Genet Elements* 2016;**6**:e1179074.
- 369 Harrison E, Guymier D, Spiers AJ *et al.* Parallel compensatory evolution stabilizes plasmids  
370 across the parasitism-mutualism continuum. *Curr Biol* 2015;**25**:2034–9.
- 371 Hartmann M, Brunner I, Hagedorn F *et al.* A decade of irrigation transforms the soil  
372 microbiome of a semi-arid pine forest. *Mol Ecol* 2017;**26**:1190–206.
- 373 Heuer H, Smalla K. Plasmids foster diversification and adaptation of bacterial populations in  
374 soil. *FEMS Microbiol Rev* 2012;**36**:1083–104.
- 375 Hughes VM, Datta N. Conjugative plasmids in bacteria of the “pre-antibiotic” era [24].  
376 *Nature* 1983;**302**:725–6.
- 377 Humeniuk C, Arlet G, Gautier V *et al.*  $\beta$ -lactamases of *Kluyvera ascorbata*, probable  
378 progenitors of some plasmid-encoded CTX-M types. *Antimicrob Agents Chemother*  
379 2002;**46**:3045–9.
- 380 Jechalke S, Heuer H, Siemens J *et al.* Fate and effects of veterinary antibiotics in soil. *Trends*  
381 *Microbiol* 2014;**22**:536–45.
- 382 Jechalke S, Kopmann C, Richter M *et al.* Plasmid-mediated fitness advantage of  
383 *Acinetobacter baylyi* in sulfadiazine-polluted soil. *FEMS Microbiol Lett* 2013;**348**:127–  
384 32.
- 385 Kennedy K, Collignon P. Colonisation with *Escherichia coli* resistant to “critically  
386 important” antibiotics: a high risk for international travellers. *Eur J Clin Microbiol*  
387 *Infect Dis* 2010;**29**:1501–6.
- 388 Klümper U, Dechesne A, Riber L *et al.* Metal stressors consistently modulate bacterial  
389 conjugal plasmid uptake potential in a phylogenetically conserved manner. *ISME J*  
390 2017;**11**:152–65.
- 391 Klümper U, Riber L, Dechesne A *et al.* Broad host range plasmids can invade an  
392 unexpectedly diverse fraction of a soil bacterial community. *ISME J* 2014;**9**:934–45.
- 393 Knapp CW, Dolfing J, Ehlert PAI *et al.* Evidence of increasing antibiotic resistance gene  
394 abundances in archived soils since 1940. *Environ Sci Technol* 2010;**44**:580–7.
- 395 Laxminarayan R, Duse A, Wattal C *et al.* Antibiotic resistance - the need for global solutions.  
396 *Lancet Infect Dis* 2013;**13**:1057–98.
- 397 Leonard AFC, Zhang L, Balfour AJ *et al.* Exposure to and colonisation by antibiotic-resistant  
398 *E. coli* in UK coastal water users: Environmental surveillance, exposure assessment, and  
399 epidemiological study (Beach Bum Survey). *Environ Int* 2018, DOI:  
400 10.1016/j.envint.2017.11.003.
- 401 Loftie-Eaton W, Bashford K, Quinn H *et al.* Compensatory mutations improve general  
402 permissiveness to antibiotic resistance plasmids. *Nat Ecol Evol* 2017;**1**:1354–63.
- 403 Lowenthal D. *The Past Is a Foreign Country – Revisited*. Cambridge: Cambridge University  
404 Press, 2015.
- 405 Mangat CS, Bekal S, Irwin RJ *et al.* A novel hybrid plasmid carrying multiple antimicrobial  
406 resistance and virulence genes in salmonella enterica serovar Dublin. *Antimicrob Agents*  
407 *Chemother* 2017;**61**:e02601-16.

1  
2  
3 408 McEwen SA, Collignon PJ. Antimicrobial resistance: a One Health perspective. *Microbiol*  
4 409 *Spectr* 2018;**6**:1–26.

5  
6 410 McKinnon J, Chowdhury PR, Djordjevic SP. Genomic analysis of multiple drug resistant  
7 411 *Escherichia coli* ST58 causing urosepsis. *Int J Antimicrob Agents* 2018;**0**:1–6.

8  
9 412 Musovic S, Klümper U, Dechesne A *et al.* Long-term manure exposure increases soil  
10 413 bacterial community potential for plasmid uptake. *Environ Microbiol Rep* 2014;**6**:125–  
11 414 30.

12  
13 415 Porse A, Schønning K, Munck C *et al.* Survival and evolution of a large multidrug resistance  
14 416 plasmid in new clinical bacterial hosts. *Mol Biol Evol* 2016;**33**:2860–73.

15  
16 417 Reid CJ, Wyrsh ER, Roy Chowdhury P *et al.* Porcine commensal *Escherichia coli*: a  
17 418 reservoir for class 1 integrons associated with IS26. *Microb Genomics* 2017;**3**:1–13.

18  
19 419 Rizzo L, Manaia C, Merlin C *et al.* Urban wastewater treatment plants as hotspots for  
20 420 antibiotic resistant bacteria and genes spread into the environment: A review. *Sci Total*  
21 421 *Environ* 2013;**447**:345–60.

22  
23 422 Robinson TP, Bu DP, Carrique-Mas J *et al.* Antibiotic resistance is the quintessential One  
24 423 Health issue. *Trans R Soc Trop Med Hyg* 2016;**110**:377–80.

25  
26 424 Ryan RP, Monchy S, Cardinale M *et al.* The versatility and adaptation of bacteria from the  
27 425 genus *Stenotrophomonas*. *Nat Rev Microbiol* 2009;**7**:514–25.

28  
29 426 San Millan A, Escudero JA, Gifford DR *et al.* Multicopy plasmids potentiate the evolution of  
30 427 antibiotic resistance in bacteria. *Nat Ecol Evol* 2016;**1**:0010.

31  
32 428 Song J, Rensing C, Holm PE *et al.* Comparison of metals and tetracycline as selective agents  
33 429 for development of tetracycline resistant bacterial communities in agricultural soil.  
34 430 *Environ Sci Technol* 2017;**51**:3040–7.

35  
36 431 Venturini C, Beatson SA, Djordjevic SP *et al.* Multiple antibiotic resistance gene recruitment  
37 432 onto the enterohemorrhagic *Escherichia coli* virulence plasmid. *FASEB J*  
38 433 2010;**24**:1160–6.

39  
40 434 Wellington EM, Boxall AB, Cross P *et al.* The role of the natural environment in the  
41 435 emergence of antibiotic resistance in Gram-negative bacteria. *Lancet Infect Dis*  
42 436 2013;**13**:155–65.

43  
44 437 Wistrand-Yuen E, Knopp M, Hjort K *et al.* Evolution of high-level resistance during low-  
45 438 level antibiotic exposure. *Nat Commun* 2018;**9**, DOI: 10.1038/s41467-018-04059-1.

46  
47 439 Wolters B, Jacquiod S, Sørensen SJ *et al.* Bulk soil and maize rhizosphere resistance genes,  
48 440 mobile genetic elements and microbial communities are differently impacted by organic  
49 441 and inorganic fertilization. *FEMS Microbiol Ecol* 2018;**94**:1–13.

50  
51 442 Wolters B, Widyasari-Mehta A, Kreuzig R *et al.* Contaminations of organic fertilizers with  
52 443 antibiotic residues, resistance genes, and mobile genetic elements mirroring antibiotic  
53 444 use in livestock? *Appl Microbiol Biotechnol* 2016;**100**:9343–53.

54  
55 445 Wong MHY, Chan EWC, Chen S. IS26-mediated formation of a virulence and resistance  
56 446 plasmid in *Salmonella* Enteritidis. *J Antimicrob Chemother* 2017;**72**:2750–4.

57  
58 447 Zhao WH, Hu ZQ. Epidemiology and genetics of CTX-M extended-spectrum  $\beta$ -lactamases in  
59 448 Gram-negative bacteria. *Crit Rev Microbiol* 2013;**39**:79–101.



- 1  
2  
3 449 Zhu YG, Gillings M, Simonet P *et al.* Microbial mass movements. *Science* 2017a;**357**:1099–  
4 450 100.  
5  
6 451 Zhu YG, Johnson TA, Su J-Q *et al.* Diverse and abundant antibiotic resistance genes in  
7 452 Chinese swine farms. *PNAS* 2013;**110**:3435–40.  
8  
9 453 Zhu YG, Zhao Y, Li B *et al.* Continental-scale pollution of estuaries with antibiotic resistance  
10 454 genes. *Nat Microbiol* 2017b;**2**:16270.  
11 455  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1     **Figure legend**

2

3     **Figure 1:** Key questions about the dynamics of antibiotic resistance. The ecology and  
4     evolution of antibiotic resistance is influenced by a series of nested interactions. Each of  
5     these can be viewed from different perspectives, summarized as the elements depicted in the  
6     diagram. The key questions that are important for understanding and managing resistance  
7     dynamics are summarized on the right hand side. Each of these questions can be applied  
8     across the different scales of the illustrated hierarchy.

9

10

For Peer Review




Figure 1: Key Questions about the dynamics of antibiotic resistance

190x254mm (96 x 96 DPI)